

# Alcohol consumption

## Frequency of binge drinking during the last year (percentage)

Sex	Age (years)	Never	< monthly	1-3 times per month	Once per week	> once per week	n
Men	18-24	4.4	21.7	40.3	22.3	11.4	1,983
	25-44	8.1	54.4	26.4	7.6	3.6	8,305
	45-64	13.8	53.0	21.1	6.9	5.3	13,157
	65+	33.3	47.8	13.3	3.0	2.5	6,130
	Total	15.6	50.2	22.3	7.3	4.6	29,575
Women	18-24	10.5	39.1	37.4	10.2	2.8	3,442
	25-44	26.1	58.7	12.6	2.1	0.6	13,950
	45-64	35.8	51.7	9.2	2.0	1.4	19,153
	65+	62.8	31.0	4.7	1.0	0.5	5,729
	Total	34.2	50.2	12.0	2.6	1.1	42,274

## Weekly alcohol consumption (units)

Sex	Age (years)	Percentiles							n
		5%	10%	25%	50%	75%	90%	95%	
Men	18-24	0	0	5	12	23	34	46	2,065
	25-44	0	0	3	6	11	20	26	8,602
	45-64	0	1	4	9	16	25	34	13,691
	65+	0	1	5	10	17	28	35	6,396
	Total	0	1	4	8	15	25	34	30,754
Women	18-24	0	0	2	6	11	17	22	3,600
	25-44	0	0	1	3	6	10	13	14,820
	45-64	0	0	2	5	9	15	20	20,175
	65+	0	0	2	5	10	16	22	6,322
	Total	0	0	2	4	8	14	18	44,917

The Danish Health Examination Survey 2007-2008 (DANHES 2007-2008) is one of the largest health examination surveys among the Danish adult population. The survey consisted of a questionnaire (n=76,484) and a health examination (n=18,065). The DANHES 2007-2008 was carried out by the National Institute of Public Health (NIPH), University of Southern Denmark, and was funded by the Ministry of the Interior and Health and the Tryg Foundation. The above results are based on non-weighted data.